

A CHECKLIST OF AQUATIC AND SEMI-AQUATIC BUGS (INSECTA:HEMIPTERA:HETEROPTERA) FROM PULAU TIOMAN, PENINSULAR MALAYSIA

C. M. Yang, Timothy C. M. Wong, H. K. Lua and L. M. Koh

Department of Biological Sciences, National University of Singapore, Kent Ridge, Singapore 119260

ABSTRACT. - A list of aquatic and semi-aquatic Hemiptera from Pulau Tioman is given with notes on their habitats. A total of 8 families, 25 genera, and 33 species are recorded from the island. *Rhagovelia sumatrensis*, *Perittopus breddini*, *Ptilomera tigrina* and *Metrocoris nigrofasciodes* are the most dominant species found in the hill streams. *Ventidius modulatus* is common both in freshwater and brackish water habitats. Three species, *Stenobates biroi*, *S. insularis* and *Hebrus nereis* are new records for Peninsular Malaysia.

KEY WORDS. - Aquatic and semi-aquatic bugs, check-list, Tioman, Malaysia.

INTRODUCTION

Two intensive studies on the fauna of Pulau Tioman, Pahang, Malaysia had been carried out by Bullock & Medway in 1966 and Day & Mowbray in 1990. Only the former survey studied the diversity of arthropod fauna on the island and paid attention to the aquatic habitats (Bullock, 1966a). The work revealed that aquatic insects were well represented, but true aquatic bugs (Nepomorpha) were virtually absent. Bullock documented members of Halovelinae, Halobatinae from the open sea and mangrove; and Gerridae, Veliidae and Hydrometridae from freshwater streams, ponds and pools. However, no species list was given. Subsequent records of aquatic bugs from the island can be found in Cheng & Fernando (1969), Fernando & Cheng (1974) and Andersen (1989). The present work provides a check-list of the aquatic heteropteran fauna of the island.

MATERIAL AND METHODS

Tioman is basically a granite island. Streams on hillsides are generally rocky and steep. Fast water flowing over large boulders forms many cascades and rock-pools which can be




Fig. 1. Map of Pulau Tioman. G = Gunung; Kg = Kampong; S = Sungei; Tg = Tanjong.

as deep as 2 metres in some areas. Vegetation is rarely found on banks or submerged in water. Headwaters are always underground, seeping through rocks and boulders. The coast is lined with large boulders and the stream waters flow quickly into the sea during low tide (e.g. Sg. Raya). In the flat coastal areas the larger streams tend to meander, such as Sg. Keliling, Sg. Baharu, Sg. Paya and Sg. Mentawak. Sg. Paya has a permanent intertidal mangrove area with sandy substrate. The river mouths of Sg. Baharu and Sg. Mentawak are sandy with coral rocks and do not have intertidal banks. A few saline pools are located at Sg. Tekek and Tg. Dungun.

The material reported here were collected in three periods and were either deposited at the Bishop Museum (BP), Honolulu or the Zoological Reference Collection (ZRC), Singapore:

- 1) 1962: 18-20 March & 10 April all collected from Kg. Tekek by K. J. Kuncheria. Material was loaned from Dr. D. A. Polhemus, Bishop Museum (BP).
- 2) 1996: 24-29 June collected by W. W. Chan, K. M. Chang, C. W. L. Fock, A. L. M. Koh, H. K. Lua, C. S. Tang, T. C. M. Wong and C. M. Yang during the 1996 Zoology Honours Field Course, National University of Singapore (ZRC).
- 3) 1997: 23-27 June unless otherwise stated, all specimens were collected by K. K. P. Lim, H. K. Lua and T. C. M. Wong during the 1997 Zoology Honours Field Course, National University of Singapore (ZRC).

The collections made in 1996 and 1997 were from various habitats at the following locations (Fig. 1) (Gg.= Gunung; Kg.= Kampung; Sg.= Sungai; Tg.= Tanjong):

Sg. Asah and Mukut waterfall (26 June 1996, 24 June 1997)
 Sg. Ayer Besar and Juara waterfall (25 June 1996, 23 June 1997)
 Sg. Baharu (28 June 1996, 27 June 1997)
 Sg. Genting (27 June 1996)
 Sg. Kelepak (26 June 1996)
 Sg. Keliling (27 June 1996, 26 June 1997)
 Sg. Mentawak (26 June 1996, 24 June 1997)
 Sg. Nipah (28 June 1996, 24 June 1997)
 Sg. Paya (25 & 27 June 1996, 25 June 1997)
 Sg. Raya (26 June 1996)
 Sg. Salang (25 June 1997)
 Sg. Tekek (24 June 1996)
 Gg. Kajang (28 June 1996) a rock stream, 500 m a.s.l.
 Kg. Tekek (25, 28 & 29 June 1996)
 Monkey Bay (25 June 1997)
 Tg. Dungun (24 June 1997)

INFRAORDER GERROMORPHA

FAMILY MESOVELIIDAE

Mesovelia horvathi Lundblad, 1933

Material examined. - 1 male, 5 nymphs, Sg. Ayer Besar. — 8 males, 6 females, Sg. Paya. — 4 males, 5 females, Sg. Keliling. — 1 male, 4 females, Sg. Dungun.

***Mesovelia vittigera* Horvath, 1895**

Material examined. - 7 males, 13 females, tidal, river mouth of Sg. Raya. — 1 female, Sg. Genting. — 1 female, Sg. Paya. — 1 male, 1 female, Sg. Keliling. — 5 males, 4 females, tidal, Sg. Kelepak.

Remarks. - Mesoveliids are commonly found among marginal vegetation on standing waters of ponds and streams. *M. vittigera* was collected from both freshwater and brackish-water habitats.

FAMILY HEBRIDAE

***Hebrus nereis* Polhemus & Polhemus, 1989**

Material examined. - 9 males, 13 females, Sg. Paya.

Remarks. - *H. nereis* is a new record for Peninsular Malaysia. Specimens were found under gravels, in holes of decaying logs submerged in brackish-water or on mangrove shore. This species was previously known from mangrove swamps in Singapore and Thailand (Polhemus & Polhemus, 1989).

***Hebrus* sp.**

Material examined. - 1 female, stagnant pond, Kg. Tekek. — 1 male, 1 female, Mukut waterfall.

***Timasius* sp.**

Material examined. - 3 males, 1 female, Juara waterfall, Sg. Ayer Besar. — 1 male, 2 females, Sg. Keliling. — 2 males, Mukut waterfall. — 1 female, Sg. Paya.

Remarks. - *Timasius* was collected from wet rocks at waterfalls and along streams. This is likely to be a new species as the buccula and metanotal are different from known species described by Andersen (1981).

FAMILY HYDROMETRIDAE

***Hydrometra jaczewskii* Lundblad, 1933**

Material examined. - 2 male, 10 females, Sg. Baharu. — 1 female, Sg. Keliling.

***Hydrometra maidli* Hungerford & Evans, 1934**

Material examined. - 1 male, 1 nymph, Sg. Ayer Besar. — 8 males, 3 females, Sg. Keliling. — 1 female, Sg. Salang.

Remarks. - Found in the weedy margins of streams, ponds and pools. *H. maidli* is the most common and widespread *Hydrometra* species in Malaysia and Singapore. It has also been

recorded from brackish waters in Singapore by Murphy (1990). Hydrometrids have been found in the stomach contents of *Rana erythraea*, a common frog collected from a stagnant pond near Kg. Tekek (Bullock, 1966b).

FAMILY VELIIDAE

SUBFAMILY MICROVELIINAE

Microvelia sp. 1

Material examined. - 2 females, Sg. Ayer Besar. — 1 female, Sg. Paya. — 5 males, 1 female, Sg. Salang. — 5 males, 6 females, Sg. Baharu.

Microvelia sp. 2

Material examined. - 1 male, 1 female, 1 macropterous female, Sg. Baharu. — 2 males, 4 females, 1 macropterous male and 1 macropterous female, Sg. Keliling.

Microvelia sp. 3

Material examined. - 1 female, 2 macropterous males, 1 macropterous female, Sg. Paya.

Remarks. - All *Microvelia* species were collected from small puddles along forest streams.

SUBFAMILY HALOVELIINAE

Halovelina abdominalis Andersen, 1989

Material examined. - 4 females, Tekek Bay — 1 nymph, Monkey Bay.

Remarks. - This marine bug is easily recognized by the upturned connexiva to the dorsum and very narrow and long abdomen. This species was previously collected from rock pools at low tide from Tioman, and at light from the Java Sea (Andersen, 1989). The four adults were collected along with *Halobates proavus*.

Haloveloides sundaensis Andersen, 1991

Material examined. - 13 nymphs, Monkey Bay, H. H. Tan.

Remarks. - The specimens were found under submerged rocks. This species has been collected from coasts of Mersing, Penang and Langkawi, and in mangroves and coastal areas of the Sunda Shelf (Andersen, 1991).

Xenobates sp. 1

Material examined. - 3 males, 2 females, tidal margins of Sg. Paya. — 44 males, 21 females, among vegetation, tidal margins of Sg. Mentawak.

Xenobates sp. 2

Material examined. - numerous males and females, tidal margins of Sg. Baharu.

Remarks. - Both species are found along the margins of very slow flowing mangrove streams. Large groups of individuals were found during low tide at Sg. Baharu, among floating vegetation near the river mouth, about 200 metres from the coast. They are not sympatric in the same estuaries on Tioman. These two undetermined *Xenobates* will be described by Andersen, who is revising the genus (Andersen, pers. comm.).

SUBFAMILY RHAGOVELIINAE

Rhagovelia sumatrensis Lundblad, 1933

Material examined. - 16 males, 17 females, 1 macropterous male, 1 macropterous female, Kg. Tekek (BP); numerous macropterous and apterous males and females, various flowing streams.

Remarks. - This is the commonest and most widespread of the five known species of *Rhagovelia* in forest streams of Peninsular Malaysia and Singapore (Yang & Polhemus, 1994). It is very abundant in slow to moderately fast-flowing streams on the island. Groups of individuals are usually found on slow riffles or behind stones on moving water.

SUBFAMILY PERITTOPINAE

Perittopus breddini Kirkaldy, 1901

Material examined. - 1 male, 1 female, Kg. Tekek (BP); numerous apterous and macropterous males and females, various forest streams and waterfalls.

Remarks. - This beautiful chestnut red bug is very common on sheltered puddles, side pools and rock pools along forest streams and waterfalls on the island. Macropterous individuals are always more numerous than apterous ones.

FAMILY GERRIDAE

SUBFAMILY RHAGADOTARSINAE

Rhagadotarsus kraepelini Breddin, 1905

Material examined. - 5 males, 12 females, brackish, Sg. Baharu. — 4 males, 7 females, 1 macropterous male, Sg. Keliling.

Remarks. - This species is common on standing waters of ponds, lakes, reservoirs, temporary pools and also quiet parts of flowing streams in Southeast Asia. It is rare on Tioman because most of the streams are fast flowing. However, a population was found on the slightly brackish region of Sg. Baharu. It was collected along with *Ventidius modulatus* and *Stenobates biroi*. Specimens of *Rhagadotarsus* and *Ventidius* were at the quieter margin of the stream, whereas *Stenobates* occurred midstream with stronger currents. It has been collected from mangroves in Singapore (Polhemus & Karunaratne, 1993) and in Sri Lanka (Andersen & Foster, 1992).

SUBFAMILY GERRINAE

Limnogonus fossarum Fabricius, 1775

Material examined. - 2 males and 2 females (all macropterous), Kg. Tekek (BP). — 2 males, Sg. Ayer Besar. — 5 males, 3 females, 2 macropterous females, Sg. Paya. — 1 macropterous female, Sg. Genting. — 12 males, 12 females, 4 macropterous males, 6 macropterous females, Sg. Keliling. — 3 males, 1 female, 1 macropterous male, 1 macropterous female, freshwater and brackish-water of Sg. Baharu.

Remarks. - This is the commonest and most widespread of Malaysian gerrids and is found in all sorts of standing waters to very slow flowing streams or even brackish waters. Macropterous forms are common, thus it can colonize temporary habitats successfully (Cheng & Fernando, 1969).

Tenagogonus (Limnometra) ciliatus Mayr, 1865

Material examined. - 1 macropterous female, brackish stagnant pool, Kg. Tekek. — 1 male, 1 female, 5 macropterous male, 2 macropterous female, tributary of Sg. Keliling. — 7 males, 3 females, 8 macropterous males, stagnant pool, Tg. Dungun. — 1 male, Sg. Keliling. — 1 macropterous female, Sg. Baharu.

Tenagogonus (Limnometra) femoratus Mayr, 1865

Material examined. - All macropterous. 1 male, 2 females, saline pool, stagnant, Kg. Tekek. — 3 females, brackish, Sg. Keliling. — 1 female, Sg. Baharu.

Remarks. - These two species of *Tenagogonus* are not common and were found in stagnant waters of both brackish pools and at quiet edges of forest streams. Although Cheng & Fernando (1969) reported that this species was only found to inhabit forest ponds, it has colonized brackish-waters on Tioman.

SUBFAMILY CYLINDROSTETHINAE

Cylindrostethus malayensis D. Polhemus, 1994

Material examined. - 1 male, Sg. Baharu.

Remarks. - This species is very rare on the island, only one specimen was collected from an open area of a stream with moderate water flow. It has previously been recorded as *C.*

SUBFAMILY ANISOPINAE

Anisops tahitiensis Lundblad, 1934

Material examined. - 8 males, 12 females, 14 nymphs, Sg. Ayer Besar. — 6 males, 4 females, 11 nymphs, Sg. Raya. — 3 males, 1 female, Sg. Keliling.

Nychia sappho Kirkaldy, 1901

Material examined. - one male, stagnant pool at Tg. Dungun.

Remarks. - It was previously known as *N. malayana* Lundblad, 1933.

FAMILY PLEIDAE

Plea quinquemaculata Lundblad, 1933

Material examined. - 6 males, 6 females, stagnant pond, Kg. Tekek. — 1 male, 3 females, stagnant pool, near Sg. Raya.

FAMILY BELOSTOMATIDAE

Diplonychus rusticus (Fabricius, 1781)

Material examined. - 10 nymphs, Sg. Ayer Besar.

DISCUSSION

Based on published literature there are about 40 genera and 100 species of Gerromorpha on the mainland of Peninsular Malaysia. The present study has documented only 25 genera and 33 species (or 60% and 33% respectively) on Pulau Tioman.

Rhagovelia sumatrensis, *Metrocoris nigrofasciodes*, *Ptilomera tigrina* and *Perittopus breddini* are the commonest species found on or along the rocky hill streams, followed by *Rheumatogonus intermedius*, *Ventidius modulatus* in some forest habitats. Small numbers of *Limnogonus fossarum*, *Tenagogonus ciliatus*, *T. femoratus* and *Rhagadotarsus kraepelini* are scattered in some areas of stagnant or slow flowing waters as well as the brackish environment. In the estuarine habitats *Ventidius modulatus*, *Stenobates insularis*, *S. biroi*, *Xenobates* spp. and *Hebrus nereis* are common. For the true marine bugs, *Halobates proavus* is common, *Halovelia abdominalis* and *Haloveloides sundaensis* are present too.

Three species are new records for the Malaysian fauna: *Stenobates biroi*, *S. insularis* and *Hebrus nereis*. The two former species were previously found only in Singapore, and the later was from Thailand and Singapore. The presence of these species from Tioman indicate that they could be expected to occur in mainland Peninsular Malaysia.

Since all species found on Tioman also occur on mainland Peninsular Malaysia, the island must have been connected to the mainland when there were major changes in the sea level which occurred at intervals during the Pleistocene (Bullock & Medway, 1966). According to Andersen (1982), species which are found on islands are most often those which are most widely distributed on the mainland and are known to colonize new temporary habitats.

The poor diversity of aquatic insects on Tioman could be due to the island's very steep granite terrain. The excessively well-drained watercourses have very limited quiet microhabitats. Aquatic vegetation, leaf litter and permanent stagnant waters are rare, therefore many common aquatic (Nepomorpha) and semi-aquatic (Gerromorpha) bugs, especially the former, are very poorly represented. Although the present list is probably not complete, we have found several species (5) of Nepomorphan bugs which were reported to be virtually absent by Bullock (1966a). Other aquatic insects, such as aquatic beetles, are also infrequently found (see Hendrich & Yang, 1998).

Through the kindness of Dr Dan Polhemus of the Bishop Museum, the first author was able to examine specimens of *Rhagovelia sumatrensis*, *Perittopus breddini*, *Limnogonus fossarum*, *Metrocoris nigrofasciodes*, *Halobates proavus* and *Timasius* sp. reported by Bullock in 1965. The other reported species, such as *Hydrometra* and *Mesovelia*, could not be found in the Bishop Museum nor in the University of Malaya collection. Since no species list was given by Bullock, we are therefore unable to compare the present hemipteran fauna with those collected in 1965. However, it is to be expected that some species may be more abundant during the 1965 survey, since freshwater swamps and more ponds were present then. Bullock (1966b) studied the food of amphibians and found that besides hydrometrids and veliids, the major item in the diet of *Rana erythraea* was gerrids. *Rana erythraea* was mostly collected from a stagnant pool near Kg. Tekek which no longer exists. Gerrids were also found in the stomach contents of another frog, *Rana cancrivora* which inhabits coastal habitats on the island.

ACKNOWLEDGEMENTS

The senior author is grateful to Dan Polhemus of Bishop Museum, for his help and the loan of material for this study, and to H. S. Yong (University Malaysia), J. L. Bullock (Leicester University), John Polhemus (Colorado Museum), Lanna Cheng (Scripps Institution), Peter Ng, N. Sivasothi and Alvin Wong for their help. Our special thanks are due to The Malaysian Department of Wildlife and National Parks (Perhilitan) for permission to study the fauna on the island; to W. W. Chan, A. L. M. Koh, C. W. L. Fock, K. M. Chang, C. S. Tang, H. H. Tan, C. Y. Goh and K. K. P. Lim for their help in collecting the materials; to Tommy Tan for logistic support; to K. K. P. Lim and S. Greasi for comments and help in the preparation of this manuscript. This study is partially supported by research grant (RP640083) to the first author from the National University of Singapore.

LITERATURE CITED

- Andersen, N. M., 1981. Semiaquatic bugs: phylogeny and classification of the Hebridae (Heteroptera: Gerromorpha) with revisions of *Timasius*, *Neotimasius* and *Hyrceanus*. *Syst. Entomol.*, **6**: 377-412.

- Andersen, N. M., 1982. *The semi-aquatic bugs (Hemiptera, Gerromorpha) phylogeny, adaptations, biography and classification*. Entomograph Vol. 3. Scandinavian Science Press Ltd, Klamperborg, Denmark. 455pp.
- Andersen, N. M., 1989. The coral bugs, genus *Halovelina* Bergroth (Hemiptera, Veliidae). II. Taxonomy of the *H. malaya*-group, cladistics, ecology, biology, and biogeography. *Entomol. Scand.*, **20**(2): 179-227.
- Andersen, N. M., 1991. A new genus of marine water-striders (Hemiptera, Veliidae) with five new species from Malesia. *Entomol. Scand.*, **22** (4): 389-404.
- Andersen, N. M. & W. A. Foster, 1992. Sea skaters of India, Sri Lanka, and the Maldives, with a new species and a revised key to Indian ocean species of *Halobates* and *Asclepios* (Hemiptera, Gerridae). *J. Nat. Hist.*, **26**: 533-534.
- Bullock, J. A., 1966a. Observation on the fauna of Pulau Tioman and Pulau Tulai: The food of the amphibians and reptiles. *Bull. Natn. Museum, Singapore*. **34**: 85-96.
- Bullock, J. A., 1966b. Observation on the fauna of Pulau Tioman and Pulau Tulai: Introductory report on the terrestrial arthropods. *Bull. Natn. Museum, Singapore*, **34**: 104-128.
- Bullock J. A. & L. Medway, 1966. Observations on the fauna of Pulau Tioman and Pulau Tulai: General introduction. *Bull. Natn. Museum, Singapore*, **34**: 1-8.
- Chen, P. P. & N. Nieser, 1993. A taxonomic revision of the oriental water strider genus *Metrocoris* Mayr (Hemiptera, Gerridae). Part I & II. *Steenstrupia*, **19**(1): 1-82.
- Cheng, L. & C. H. Fernando, 1969. A taxonomic study of the Malayan Gerridae (Hemiptera, Heteroptera) with notes on their biology and distribution. *Orient. Ins.*, **3**(2): 97-160.
- Day, M. & T. Mowbray, 1990. University of Bristol Tioman Archipelago Expedition, Peninsula Malaysia - Final Report. Unpublished. 90pp.
- Esaki, T., 1926. The water-striders of the subfamily Halobatinae in the Hungarian National Museum. *Ann. Mus. Nat. Hung.*, **23**: 117-164.
- Fernando, C. H. & L. Cheng, 1974. A preliminary study on the fauna of aquatic Hemiptera in Malaya and Singapore. *Fed. Mus. J.*, **19**: 21-44.
- Hendrich, L. & C. M. Yang, 1998. A contribution to the knowledge of the water beetle fauna of Pulau Tioman, Peninsular Malaysia (Coleoptera: Noteridae, Dytiscidae, Hydrophilidae, Hydraenidae, Scirtidae, Limnichidae). *Raffles Bull. Zool.*, **46**(2): 1-10.
- Murphy, D. H., 1990. Walkers on water - An account of the pleuston of Singapore. In: Chou L. M. & P. K. L. Ng (eds) Pp. 153-168 *Essays in Zoology*. Department of Zoology National University of Singapore.
- Polhemus, D. A., 1994. Taxonomy, phylogeny, and zoogeography of the genus *Cylindrostethus* Fiber in the Palearctic region (Heteroptera: Gerridae). *Bishop Mus. Occ. Pap.*, **38**: 1-34.
- Polhemus, J. T. & L. Cheng, 1982. Notes on marine water-striders with descriptions of new species. *Pacific Ins.*, **24** (3-4): 219-227.
- Polhemus, J. T. & R. P. Karunaratne, 1993. A review of the genus *Rhagadotarsus*, with description of three new species (Heteroptera: Gerridae). *Raffles Bull. Zool.*, **41**(1): 97-114.
- Polhemus, J. T. & D. A. Polhemus, 1989. A new mesoveliid genus and two new species of *Hebrus* (Heteroptera: Mesoveliidae, Hebridae) from intertidal habitats in Southeast Asian mangrove swamps. *Raffles Bull. Zool.*, **37** (1& 2): 73-82.
- Polhemus, J. T. & D. A. Polhemus, 1991. Three new species of marine water-striders from the Australasian region, with notes on other species (Gerridae: Halobatinae, Trepobatinae). *Raffles Bull. Zool.*, **39**(1): 1-14.
- Yang, C. M. & D. Kovac, 1995. A collection of aquatic and semi-aquatic bugs (Insecta :Hemiptera: Gerromorpha and Nepomorpha) from Tememgor Forest Reserve, Hulu Perak, Malaysia. *Malayan Nat. J.*, **48**: 287-296.
- Yang, C. M. & D. A. Polhemus, 1994. Notes on *Rhagovelia* Mayr (Hemiptera: Veliidae) from Singapore, with description of a new species. *Raffles Bull. Zool.*, **42**(2): 987-993.